

OXIDIZING COMPOSITION AND USES FOR DYEING, FOR
PERMANENTLY RESHAPING OR FOR BLEACHING KERATIN FIBRES

The present application relates to a cosmetic composition intended for treating keratin fibres, comprising, in a support which is suitable for keratin fibres:

(a) at least one enzyme of 2-electron oxidoreductase type in the presence of at least one donor for the said enzyme;

(b) at least one aminosilicone;

as well as to processes for treating keratin fibres, in particular processes for dyeing, permanently reshaping or bleaching the hair, using this composition.

00832877 044204

OXIDIZING COMPOSITION AND USES FOR DYEING, FOR
PERMANENTLY RESHAPING OR FOR BLEACHING KERATIN FIBRES

The present invention relates to an oxidizing
5 composition intended for treating keratin fibres,
comprising at least one enzyme of 2-electron
oxidoreductase type in the presence of at least one
donor for the said enzyme and at least one
aminosilicone as well as to its uses for dyeing, for
10 permanently reshaping or for bleaching keratin fibres,
in particular human hair.

It is known to dye keratin fibres, and in
particular human hair, with dye compositions containing
oxidation dye precursors, in particular para-
15 phenylenediamines, ortho- or para-aminophenols and
heterocyclic bases which are generally referred to as
oxidation bases. Oxidation dye precursors, or oxidation
bases, are colourless or weakly coloured compounds
which, when combined with oxidizing products, can give
20 rise to coloured compounds and dyes by a process of
oxidative condensation.

It is also known that the shades obtained with
these oxidation bases can be varied by combining them
with couplers or colour modifiers, the latter being
25 chosen in particular from aromatic meta-diamines, meta-
aminophenols, meta-diphenols and certain heterocyclic
compounds.

The variety of compounds used as regards the
oxidation bases and the couplers allows a wide range of
30 colours to be obtained.

The so-called "permanent" coloration obtained
by means of these oxidation dyes must moreover satisfy
a certain number of requirements. Thus it must have no
toxicological drawbacks, it must be able to give shades
35 of the desired intensity and it must be able to
withstand external agents (light, bad weather, washing,

0983287.04101

permanent-waving, perspiration, rubbing).

The dyes must also be able to cover white hair and, lastly, they must be as unselective as possible, i.e. they must give the smallest possible colour differences along the same length of keratin fibre, which may in fact be differently sensitized (i.e. damaged) between its tip and its root.

The oxidation dyeing of keratin fibres is generally carried out in alkaline medium, in the presence of hydrogen peroxide. However, the use of alkaline media in the presence of hydrogen peroxide has the drawback of causing appreciable degradation of the fibres, as well as considerable bleaching of the keratin fibres, which is not always desirable.

The oxidation dyeing of keratin fibres can also be carried out using oxidizing systems other than hydrogen peroxide, such as enzymatic systems. Thus, it has already been proposed to dye keratin fibres, in particular in patent application EP-A-0,310,675, with compositions comprising an oxidation dye precursor in combination with enzymes such as pyranose oxidase, glucose oxidase or uricase, in the presence of a donor for the said enzymes. Although being used under conditions which do not result in degradation of the keratin fibres which is comparable to that caused by the dyes used in the presence of hydrogen peroxide, these dye formulations nevertheless lead to colorations which are still insufficient, both as regards the homogeneity of the colour distributed along the fibre ("unison") and as regards the chromaticity (luminosity), the dyeing power and the resistance to the various aggressive factors to which the hair may be subjected.

It is known that the most common technique for obtaining a permanent reshaping of the hair consists, in a first stage, in opening the keratin -S-S-disulphide

(cysteine) bonds using a composition containing a suitable reducing agent (reduction step) followed, after having rinsed the hair thus treated, by reconstituting, in a second stage, the said disulphide bonds by applying to the hair, which has been placed under tension beforehand (curlers and the like), an oxidizing composition (oxidation step, also known as the fixing step) so as finally to give to the hair the desired shape. This technique thus makes it equally possible either to make the hair wavy or to straighten it or to remove its curliness. The new shape given to the hair by a chemical treatment such as above is remarkably long-lasting and in particular resists the action of washing with water or shampoos, as opposed to simple standard techniques for temporary reshaping, such as hairsetting.

The reducing compositions which may be used in order to carry out the first step of a permanent-waving operation generally contain, as reducing agents, sulphites, bisulphites, alkylphosphines or, preferably, thiols. Among the thiols, those commonly used are cysteine and the various derivatives thereof, cysteamine and the derivatives thereof, thiolactic acid or thioglycolic acid, the salts thereof and the esters thereof, in particular glyceryl thioglycolate.

As regards the oxidizing compositions needed to carry out the fixing step, use is usually made in practice of compositions based on aqueous hydrogen peroxide, sodium bromate or persalts such as sodium perborate, which have the drawback of being liable to damage the hair.

The problem of the technique of the permanent-waving operations known to date is that their application to the hair induces long-term adverse changes in the quality of the hair. The essential causes of these adverse changes in the quality of the hair are a reduction in its cosmetic properties, such

00832877 044201

as its sheen and its feel, and degradation of its mechanical properties, more particularly degradation of its mechanical strength due to swelling of the keratin fibres during the rinsing between the reduction step
5 and the oxidation step, which can also be reflected by an increase in its porosity. The hair is weakened and can become brittle during subsequent treatments such as blow-drying.

10 The same problem of adverse changes in keratin fibres is encountered during processes for bleaching the hair.

It is known that the permanent reshaping or bleaching of keratin fibres can also be carried out under milder conditions using oxidizing systems other
15 than hydrogen peroxide, such as enzymatic systems. Thus, processes for the permanent reshaping or bleaching of keratin fibres have already been proposed, in particular in patent application EP-A-0,310,675, with compositions comprising an enzyme such as pyranose
20 oxidase, glucose oxidase or uricase, in the presence of a donor for the said enzyme. Although being used under conditions which do not result in degradation of the keratin fibres which is comparable to that caused by conventional permanent-waving or bleaching processes,
25 these oxidizing formulations nevertheless lead to results which are still insufficient, as regards the curl hold over time, as regards the compatibility of permanent-waved or bleached hair with subsequent treatments, as regards the degradation of the
30 mechanical properties of the permanent-waved hair, in particular the reduction of the porosity of the hair, and as regards the reduction of the cosmetic properties such as the feel, or alternatively as regards the uniformity of the bleaching along the keratin fibres.

35 The aim of the present invention is to solve the problems mentioned above.

composition gives a more uniform bleaching effect on the hair and improves the cosmetic properties, such as the feel.

These discoveries form the basis of the present invention.

The subject of the present invention is thus, firstly, a cosmetic and/or dermatological composition intended for treating keratin fibres, in particular human keratin fibres and more particularly human hair, comprising, in a support which is suitable for keratin fibres:

- (a) at least one enzyme of 2-electron oxidoreductase type in the presence of at least one donor for the said enzyme,
- (b) at least one aminosilicone.

The 2-electron oxidoreductase(s) used in the oxidizing compositions in accordance with the invention can be chosen in particular from pyranose oxidases, glucose oxidases, glycerol oxidases, lactate oxidases, pyruvate oxidases and uricases.

According to the invention, the 2-electron oxidoreductase is preferably chosen from uricases of animal, microbiological or biotechnological origin.

By way of example, mention may be made in particular of uricase extracted from boar liver, uricase from *Arthrobacter globiformis*, as well as uricase from *Aspergillus flavus*.

The 2-electron oxidoreductase(s) can be used in pure crystalline form or in a form diluted in a diluent which is inert with respect to the said 2-electron oxidoreductase.

The 2-electron oxidoreductase(s) in accordance with the invention preferably represent(s) from 0.01 to 20% by weight approximately relative to the total weight of the composition, and even more preferably from 0.1 to 5% by weight approximately relative to this

weight.


According to the invention, the term donor is understood to refer to the various substrates also necessary for the functioning of the said 2-electron oxidoreductase(s).

The nature of the donor (or substrate) for the said enzyme varies depending on the nature of the 2-electron oxidoreductase used. For example, as donors for the pyranose oxidases, mention may be made of D-glucose, L-sorbose and D-xylose; as a donor for the glucose oxidases, mention may be made of D-glucose; as donors for the glycerol oxidases, mention may be made of glycerol and dihydroxyacetone; as donors for the lactate oxidases, mention may be made of lactic acid and its salts; as donors for the pyruvate oxidases, mention may be made of pyruvic acid and its salts; and lastly, as donors for the uricases, mention may be made of uric acid and its salts.

The donor(s) (or substrate(s)) used in accordance with the invention preferably represent(s) from 0.01 to 20% by weight approximately relative to the total weight of the composition in accordance with the invention, and even more preferably from 0.1 to 5% approximately relative to this weight.


According to the invention, the term aminosilicone is understood to denote any silicone containing at least one primary, secondary or tertiary amine or a quaternary ammonium group. Mention may thus be made of:

(a) the polysiloxanes referred to in the CTFA dictionary as "amodimethicone" and corresponding to formula (I):



in which x and y are integers dependent on the molecular weight, generally such that the said number-average molecular weight is between 5000 and 500,000 approximately;

(b) cationic silicone polymers corresponding to the formula:



in which:





G is a hydrogen atom or a phenyl, OH or C₁-C₈ alkyl, for example methyl, group,



a denotes the number 0 or an integer from 1 to 3, in particular 0,

b denotes 0 or 1, and in particular 1,

m and n are numbers such that the sum (n + m) can range especially from 1 to 2000 and in particular from 50 to 150, it being possible for n to denote a number from 0 to 1999 and in particular from 49 to 149, and it being possible for m to denote a number from 1 to 2000 and in particular from 1 to 10;


R is a monovalent radical of formula -C_qH_{2q}L in which q is a number from 2 to 8 and L is an optionally quaternized amine group chosen from the groups:






in which R' can denote hydrogen, phenyl, benzyl or a monovalent, saturated hydrocarbon-based radical, for example an alkyl radical containing from 1 to 20 carbon atoms, and A⁻ represents a halide ion such as, for example, fluoride, chloride, bromide or iodide.

A product corresponding to this definition is the polymer known as "trimethylsilylamodimethicone" corresponding to formula (III) below:



in which n and m have the meanings given above for formula (II). Such polymers are described, for example, in patent application EP-A-95238.

(c) cationic silicone polymers corresponding to the formula:



in which

R₁ represents a monovalent hydrocarbon-based radical containing from 1 to 18 carbon atoms, and in particular a C₁-C₁₈ alkyl radical or a C₂-C₁₈ alkenyl radical, for example methyl;

R₂ represents a divalent hydrocarbon-based radical, in particular a C₁-C₁₈ alkylene radical or a C₁-C₁₈, for example C₁-C₈, divalent alkyleneoxy radical;

Q⁻ is a halide ion, in particular chloride;

r represents an average statistical value from 2 to 20 and in particular from 2 to 8;


s represents an average statistical value from 20 to 200 and in particular from 20 to 50.

Such polymers are described more particularly in US patent 4,185,087.

One polymer belonging to this class is the polymer sold by the company Union Carbide under the name "Ucar Silicone ALE 56".

When these silicone polymers are used, one particularly advantageous embodiment is their use together with cationic and/or nonionic surfactants.

It is possible, for example, to use the product sold under the name "Cationic Emulsion DC 929" by the company Dow Corning, which comprises, besides amodimethicone, a cationic surfactant comprising a mixture of products corresponding to the formula:



in which R₃ denotes alkenyl and/or alkyl radicals containing from 14 to 22 carbon atoms, derived from tallow fatty acids, and known under the CTFA name "tallowtrimonium chloride", in combination with a

nonionic surfactant of formula: $C_9H_{19}-C_6H_4-(OC_2H_4)_{10}-OH$, known under the name "Nonoxynol 10".

Use may also be made, for example, of the product sold under the name "Cationic Emulsion DC 939" by the company Dow Corning, which comprises, besides amodimethicone, a cationic surfactant which is trimethylcetylammonium chloride, in combination with a nonionic surfactant of formula $C_{13}H_{27}-(OC_2H_4)_{12}-OH$, known under the CTFA name "trideceth-12".

Another commercial product which can be used according to the invention is the product sold under the name "Dow Corning Q2 7224" by the company Dow Corning, containing, in combination, trimethylsilylamodimethicone of formula (III) described above, a nonionic surfactant of formula: $C_8H_{17}-C_6H_4-(OCH_2CH_2)_{40}-OH$, known under the CTFA name "octoxynol-40", a second nonionic surfactant of formula: $C_{12}H_{25}-(OCH_2-CH_2)_6-OH$, known under the CTFA name "isolaureth-6", and propylene glycol.


The compositions in accordance with the invention contain the aminosilicones defined above at weight contents which can be between 0.05% and 10%, preferably between 0.1% and 5% and even more preferably between 0.2% and 3%, relative to the total weight of the composition.

A subject of the present invention is also a ready-to-use composition for the oxidation dyeing of keratin fibres, and in particular human keratin fibres such as the hair, of the type comprising, in a medium which is suitable for dyeing, at least one oxidation base and, where appropriate, one or more couplers, which is characterized in that it contains:

- (a) at least one enzyme of 2-electron oxidoreductase type in the presence of at least one donor for the said enzyme,
- (b) at least one aminosilicone.

The nature of the oxidation base(s) used in the ready-to-use dye composition is not a critical factor. They can be chosen, in particular, from para-phenylenediamines, double bases, para-aminophenols, ortho-aminophenols and heterocyclic oxidation bases.

Among the para-phenylenediamines which can be used as oxidation bases in the dye compositions in accordance with the invention, mention may be made in particular of the compounds of formula (V) below, and the addition salts thereof with an acid:



in which:

- R_4 represents a hydrogen atom, a $\text{C}_1\text{-C}_4$ alkyl radical, a $\text{C}_1\text{-C}_4$ monohydroxyalkyl radical, a $\text{C}_2\text{-C}_4$ polyhydroxyalkyl radical, a $(\text{C}_1\text{-C}_4)\text{alkoxy}(\text{C}_1\text{-C}_4)\text{alkyl}$ radical, a $\text{C}_1\text{-C}_4$ alkyl radical substituted with a nitrogenous group, a phenyl radical or a 4'-aminophenyl radical;
- R_5 represents a hydrogen atom, a $\text{C}_1\text{-C}_4$ alkyl radical, a $\text{C}_1\text{-C}_4$ monohydroxyalkyl radical, a $\text{C}_2\text{-C}_4$ polyhydroxyalkyl radical, a $(\text{C}_1\text{-C}_4)\text{alkoxy}(\text{C}_1\text{-C}_4)\text{alkyl}$ radical or a $\text{C}_1\text{-C}_4$ alkyl radical substituted with a nitrogenous group;
- R_6 represents a hydrogen atom, a halogen atom such as a chlorine, bromine, iodine or fluorine atom, a $\text{C}_1\text{-C}_4$ alkyl radical, a $\text{C}_1\text{-C}_4$ monohydroxyalkyl radical, a $\text{C}_1\text{-C}_4$ hydroxyalkoxy radical, an acetyl-amino $(\text{C}_1\text{-C}_4)\text{alkoxy}$ radical, a $\text{C}_1\text{-C}_4$ mesylaminoalkoxy radical or a carbamoylamino $(\text{C}_1\text{-C}_4)\text{alkoxy}$ radical,

- R₇ represents a hydrogen or halogen atom or a C₁-C₄ alkyl radical.

Among the nitrogenous groups of formula (V) above, mention may be made in particular of amino, 5 mono(C₁-C₄)alkylamino, di(C₁-C₄)alkylamino, tri(C₁-C₄)alkylamino, monohydroxy(C₁-C₄)alkylamino, imidazolinium and ammonium radicals.


Among the para-phenylenediamines of formula (V) above, mention may be made more particularly of para-phenylenediamine, para-toluylenediamine, 2-chloro-para-phenylenediamine, 2,3-dimethyl-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-phenylenediamine, 2,5-dimethyl-para-phenylenediamine, N,N-dimethyl-para-phenylenediamine, N,N-diethyl-para-phenylenediamine, N,N-dipropyl-para-phenylenediamine, 15 4-amino-N,N-diethyl-3-methylaniline, N,N-bis(β-hydroxyethyl)-para-phenylenediamine, 4-amino-N,N-bis(β-hydroxyethyl)-2-methylaniline, 4-amino-2-chloro-N,N-bis(β-hydroxyethyl)aniline, 2-β-hydroxyethyl-para-phenylenediamine, 2-fluoro-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, N-(β-hydroxypropyl)-para-phenylenediamine, 2-hydroxymethyl-para-phenylenediamine, N,N-dimethyl-3-methyl-para-phenylenediamine, N,N-(ethyl-β-hydroxyethyl)-para-phenylenediamine, N-(β,γ-dihydroxypropyl)-para-phenylenediamine, N-(4'-aminophenyl)-para-phenylene- 25 diamine, N-phenyl-para-phenylenediamine, 2-β-hydroxyethyloxy-para-phenylenediamine, 2-β-acetylaminoethyloxy-para-phenylenediamine and N-(β-methoxyethyl)-para-phenylenediamine, and the addition salts thereof 30 with an acid.

Among the para-phenylenediamines of formula (V) above, para-phenylenediamine, para-toluylenediamine, 2-isopropyl-para-phenylenediamine, 2-β-hydroxyethyl-para-phenylenediamine, 35 2-β-hydroxyethyloxy-para-

phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-phenylenediamine, 2,3-dimethyl-para-phenylenediamine, N,N-bis(β-hydroxyethyl)-para-phenylenediamine, 2-chloro-para-phenylenediamine and 2-β-acetylaminoethyloxy-para-phenylenediamine and the addition salts thereof with an acid are most particularly preferred.

According to the invention, the term double bases is understood to refer to the compounds containing at least two aromatic rings bearing amino and/or hydroxyl groups.

Among the double bases which can be used as oxidation bases in the dye compositions in accordance with the invention, mention may be made in particular of the compounds corresponding to formula (VI) below, and the addition salts thereof with an acid:



20

in which:

- Z₁ and Z₂, which may be identical or different, represent a hydroxyl or -NH₂ radical which may be substituted with a C₁-C₄ alkyl radical or with a linker arm Y;
- the linker arm Y represents a linear or branched alkylene chain containing from 1 to 14 carbon atoms, which may be interrupted by or terminated with one or more nitrogenous groups and/or one or more hetero atoms such as oxygen, sulphur or nitrogen atoms, and

30

0983227-041100

optionally substituted with one or more hydroxyl or C₁-C₆ alkoxy radicals;

- R₈ and R₉ represent a hydrogen or halogen atom, a C₁-C₄ alkyl radical, a C₁-C₄ monohydroxyalkyl radical, a

5 C₂-C₄ polyhydroxyalkyl radical, a C₁-C₄ aminoalkyl radical or a linker arm Y;

- R₁₀, R₁₁, R₁₂, R₁₃, R₁₄ and R₁₅, which may be identical or different, represent a hydrogen atom, a linker arm Y or a C₁-C₄ alkyl radical;

10 it being understood that the compounds of formula (VI) contain only one linker arm Y per molecule.


Among the nitrogenous groups of formula (VI) above, mention may be made in particular of amino, mono(C₁-C₄)alkylamino, di(C₁-C₄)alkylamino, tri(C₁-
15 C₄)alkylamino, monohydroxy(C₁-C₄)alkylamino, imidazolium and ammonium radicals.

Among the double bases of formula (VI) above, mention may be made more particularly of N,N'-bis(β-hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol,
20 N,N'-bis(β-hydroxyethyl)-N,N'-bis(4'-aminophenyl)ethylenediamine, N,N'-bis(4-aminophenyl)-tetramethylenediamine, N,N'-bis(β-hydroxyethyl)-N,N'-bis(4-aminophenyl)tetramethylenediamine, N,N'-bis(4-methylaminophenyl)tetramethylenediamine, N,N'-bis(ethyl)-N,N'-bis(4'-amino-3'-methylphenyl)ethylenedi-
25 amine and 1,8-bis(2,5-diaminophenoxy)-3,5-dioxaoctane, and the addition salts thereof with an acid.

Among these double bases of formula (VI), N,N'-bis(β-hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol and 1,8-bis(2,5-diaminophenoxy)-3,5-dioxaoctane, or one of the addition salts thereof with an acid, are particularly preferred.

Among the para-aminophenols which can be used as oxidation bases in the dye compositions in
35 accordance with the invention, mention may be made in particular of the compounds corresponding to formula

(VII) below, and the addition salts thereof with an acid:



5

in which:

- R₁₆ represents a hydrogen or halogen atom or a C₁-C₄ alkyl, C₁-C₄ monohydroxyalkyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, C₁-C₄ aminoalkyl or hydroxy(C₁-C₄)alkylamino-(C₁-C₄)alkyl radical,

- R₁₇ represents a hydrogen or halogen atom or a C₁-C₄ alkyl, C₁-C₄ monohydroxyalkyl, C₂-C₄ polyhydroxyalkyl, C₁-C₄ aminoalkyl, C₁-C₄ cyanoalkyl or (C₁-C₄)alkoxy-(C₁-C₄)alkyl radical,

15 it being understood that at least one of the radicals R₁₆ or R₁₇ represents a hydrogen atom.

Among the para-aminophenols of formula (VII) above, mention may be made more particularly of para-aminophenol, 4-amino-3-methylphenol, 4-amino-3-fluorophenol, 4-amino-3-hydroxymethylphenol, 4-amino-2-methylphenol, 4-amino-2-hydroxymethylphenol, 4-amino-2-methoxymethylphenol, 4-amino-2-aminomethylphenol, 4-amino-2-(β-hydroxyethylaminomethyl)phenol and 4-amino-2-fluorophenol, and the addition salts thereof with an acid.

Among the ortho-aminophenols which can be used as oxidation bases in the dye compositions in accordance with the invention, mention may be made more particularly of 2-aminophenol, 2-amino-5-methylphenol, 2-amino-6-methylphenol and 5-acetamido-2-aminophenol, and the addition salts thereof with an acid.

Among the heterocyclic bases which can be used as oxidation bases in the dye compositions in accordance with the invention, mention may be made more particularly of pyridine derivatives, pyrimidine derivatives, pyrazole derivatives and pyrazolo-pyrimidine derivatives, and the addition salts thereof with an acid.


Among the pyridine derivatives, mention may be made more particularly of the compounds described, for example, in patents GB 1,026,978 and GB 1,153,196, such as 2,5-diaminopyridine, 2-(4-methoxyphenyl)amino-3-aminopyridine, 2,3-diamino-6-methoxypyridine, 2-(β -methoxyethyl)amino-3-amino-6-methoxypyridine and 3,4-diaminopyridine, and the addition salts thereof with an acid.

Among the pyrimidine derivatives, mention may be made more particularly of the compounds described, for example, in German patent DE 2,359,399 or Japanese patent JP 88-169,571 or patent application WO 96/15765, such as 2,4,5,6-tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 2-hydroxy-4,5,6-triaminopyrimidine, 2,4-dihydroxy-5,6-diaminopyrimidine and 2,5,6-triaminopyrimidine, and the addition salts thereof with an acid.

Among the pyrazole derivatives, mention may be made more particularly of the compounds described in patents DE 3,843,892, DE 4,133,957 and patent applications WO 94/08969, WO 94/08970, FR-A-2,733,749 and DE 195 43 988, such as 4,5-diamino-1-methylpyrazole, 3,4-diaminopyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, 4,5-diamino-1,3-dimethylpyrazole, 4,5-diamino-3-methyl-1-phenylpyrazole, 4,5-diamino-1-methyl-3-phenylpyrazole, 4-amino-1,3-dimethyl-5-hydrazinopyrazole, 1-benzyl-4,5-diamino-3-methylpyrazole, 4,5-diamino-3-tert-butyl-1-methylpyrazole, 4,5-diamino-1-tert-butyl-3-methylpyrazole,

4,5-diamino-1-(β -hydroxyethyl)-3-methylpyrazole, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-1-ethyl-3-(4'-methoxyphenyl)pyrazole, 4,5-diamino-1-ethyl-3-hydroxymethylpyrazole, 4,5-diamino-3-hydroxymethyl-1-methylpyrazole, 4,5-diamino-3-hydroxymethyl-1-isopropylpyrazole, 4,5-diamino-3-methyl-1-isopropylpyrazole, 4-amino-5-(2'-aminoethyl)amino-1,3-dimethylpyrazole, 3,4,5-triaminopyrazole, 1-methyl-3,4,5-triaminopyrazole, 3,5-diamino-1-methyl-4-methylamino-pyrazole and 3,5-diamino-4-(β -hydroxyethyl)amino-1-methylpyrazole, and the addition salts thereof with an acid.

Among the pyrazolopyrimidine derivatives, mention may be made more particularly of the pyrazolo[1,5-a]pyrimidines of formula (VIII) below, and the addition salts thereof with an acid or with a base and the tautomeric forms thereof, when a tautomeric equilibrium exists:



in which:

- R_{18} , R_{19} , R_{20} and R_{21} , which may be identical or different, denote a hydrogen atom, a C_1 - C_4 alkyl radical, an aryl radical, a C_1 - C_4 hydroxyalkyl radical, a C_2 - C_4 polyhydroxyalkyl radical, a $(C_1$ - C_4)alkoxy(C_1 - C_4)alkyl radical, a C_1 - C_4 aminoalkyl radical (it being possible for the amine to be protected with an acetyl, ureido or sulphonyl radical), a $(C_1$ - C_4)alkylamino(C_1 - C_4)alkyl radical, a di[(C_1 - C_4)alkyl]amino(C_1 - C_4)alkyl radical (it being possible for the dialkyl radicals to form a 5- or 6-membered carbon-based ring or a

09832877-041201

heterocycle), a hydroxy(C₁-C₄)alkyl- or di[hydroxy-(C₁-C₄)alkyl]amino(C₁-C₄)alkyl radical;

- the radicals X, which may be identical or different, denote a hydrogen atom, a C₁-C₄ alkyl radical, an aryl radical, a C₁-C₄ hydroxyalkyl radical, a C₂-C₄ polyhydroxyalkyl radical, a C₁-C₄ aminoalkyl radical, a (C₁-C₄)alkylamino(C₁-C₄)alkyl radical, a di[(C₁-C₄)alkyl]amino(C₁-C₄)alkyl radical (it being possible for the dialkyls to form a 5- or 6-membered carbon-based ring or a heterocycle), a hydroxy(C₁-C₄)alkyl- or di[hydroxy(C₁-C₄)alkyl]amino(C₁-C₄)alkyl radical, an amino radical, a (C₁-C₄)alkyl- or di[(C₁-C₄)alkyl]amino radical; a halogen atom, a carboxylic acid group or a sulphonic acid group;

- i is equal to 0, 1, 2 or 3;

- p is equal to 0 or 1;

- q is equal to 0 or 1;

- n is equal to 0 or 1;

with the proviso that:


- the sum p + q is other than 0;

- when p + q is equal to 2, then n is equal to 0 and the groups NR₁₈R₁₉ and NR₂₀R₂₁ occupy the (2,3); (5,6); (6,7); (3,5) or (3,7) positions;

- when p + q is equal to 1, then n is equal to 1 and the group NR₁₈R₁₉ (or NR₂₀R₂₁) and the OH group occupy the (2,3); (5,6); (6,7); (3,5) or (3,7) positions.

When the pyrazolo[1,5-a]pyrimidines of formula (VIII) above are such that they contain a hydroxyl group on one of the positions 2, 5 or 7 α to a nitrogen atom, a tautomeric equilibrium exists represented, for example, by the following scheme:

00832877-041201



Among the pyrazolo[1,5-a]pyrimidines of formula (VIII) above, mention may be made in particular of:

- 5 - pyrazolo[1,5-a]pyrimidine-3,7-diamine;
- 2,5-dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine;
- pyrazolo[1,5-a]pyrimidine-3,5-diamine;
- 2,7-dimethylpyrazolo[1,5-a]pyrimidine-3,5-diamine;
- 3-aminopyrazolo[1,5-a]pyrimidin-7-ol;
- 10 - 3-aminopyrazolo[1,5-a]pyrimidin-5-ol;
- 2-(3-aminopyrazolo[1,5-a]pyrimidin-7-ylamino)ethanol;
- 2-(7-aminopyrazolo[1,5-a]pyrimidin-3-ylamino)ethanol;
- 2-[(3-aminopyrazolo[1,5-a]pyrimidin-7-yl)-(2-hydroxyethyl)amino]ethanol;
- 15 - 2-[(7-aminopyrazolo[1,5-a]pyrimidin-3-yl)-(2-hydroxyethyl)amino]ethanol;
- 5,6-dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine;
- 2,6-dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine;
- 2,5,N7,N7-tetramethylpyrazolo[1,5-a]pyrimidine-3,7-
- 20 diamine;

and the addition salts thereof and the tautomeric forms thereof, when a tautomeric equilibrium exists.

The pyrazolo[1,5-a]pyrimidines of formula (VIII) above can be prepared by cyclization starting
25 with an aminopyrazole, according to the syntheses described in the following references:

- EP 628559 Beiersdorf-Lilly.
- R. Vishdu, H. Navedul, Indian J. Chem., 34b (6), 514, 1995.
- 30 - N.S. Ibrahim, K.U. Sadek, F.A. Abdel-Al, Arch. Pharm., 320, 240, 1987.
- R.H. Springer, M.B. Scholten, D.E. O'Brien,

T. Novinson, J.P. Miller, R.K. Robins, J. Med. Chem., 25, 235, 1982.

- T. Novinson, R.K. Robins, T.R. Matthews, J. Med. Chem., 20, 296, 1977.

5 - US 3907799 ICN Pharmaceuticals.

The pyrazolo[1,5-a]pyrimidines of formula (VIII) above can also be prepared by cyclization starting from hydrazine, according to the syntheses described in the following references:

10 - A. McKillop and R.J. Kobilecki, Heterocycles, 6(9), 1355, 1977.

- E. Alcade, J. De Mendoza, J.M. Marcia-Marquina, C. Almera, J. Elguero, J. Heterocyclic Chem., 11(3), 423, 1974.

15 - K. Saito, I. Hori, M. Higashashi, H. Midorikawa, Bull. Chem. Soc. Japan, 47(2), 476, 1974.

The oxidation base(s) in accordance with the invention preferably represent(s) from 0.0005 to 12% by weight approximately relative to the total weight of the ready-to-use dye composition, and even more preferably from 0.005 to 6% by weight approximately relative to this weight.

The couplers which can be used are those used conventionally in oxidation dye compositions, i.e. meta-phenylenediamines, meta-aminophenols and meta-diphenols, mono- or polyhydroxylated naphthalene derivatives, sesamol and its derivatives and heterocyclic compounds such as, for example, indole derivatives, indoline derivatives, benzimidazole derivatives, benzomorpholine derivatives, sesamol derivatives, pyrazoloazole derivatives, pyrroloazole derivatives, imidazoloazole derivatives, pyrazolo-pyrimidine derivatives, pyrazoline-3,5-dione derivatives, pyrrolo[3,2-d]oxazole derivatives, pyrazolo[3,4-d]thiazole derivatives, thiazoloazole S-oxide derivatives and thiazoloazole S,S-dioxide derivatives,

and the addition salts thereof with an acid.

These couplers can be chosen in particular from
2-methyl-5-aminophenol, 5-N-(β -hydroxyethyl)amino-2-
methylphenol, 3-aminophenol, 1,3-dihydroxybenzene, 1,3-
5 dihydroxy-2-methylbenzene, 4-chloro-1,3-dihydroxy-
benzene, 2,4-diamino-1-(β -hydroxyethyloxy)benzene,
2-amino-4-(β -hydroxyethylamino)-1-methoxybenzene, 1,3-
diaminobenzene, 1,3-bis(2,4-diaminophenoxy)propane,
sesamol, α -naphthol, 6-hydroxyindole, 4-hydroxyindole,
10 4-hydroxy-N-methylindole, 6-hydroxyindoline, 2,6-
dihydroxy-4-methylpyridine, 1H-3-methylpyrazol-5-one
and 1-phenyl-3-methylpyrazol-5-one, and the addition
salts thereof with an acid.

When they are present, these couplers prefer-
15 ably represent from 0.0001 to 10% by weight approxi-
mately relative to the total weight of the ready-to-use
dye composition, and even more preferably from 0.005 to
5% by weight approximately relative to this weight.

In general, the addition salts with an acid
20 which can be used in the context of the dye com-
positions of the invention (oxidation bases and
couplers) are chosen in particular from the
hydrochlorides, hydrobromides, sulphates, tartrates,
lactates and acetates.

The dye composition of the invention can also
25 contain, in addition to the oxidation dye precursors
defined above and the optional combined couplers,
direct dyes to enrich the shades with glints. These
direct dyes can then be chosen in particular from nitro
30 dyes, azo dyes or anthraquinone dyes.

The subject of the invention is also a process
for dyeing keratin fibres, and in particular human
keratin fibres such as the hair, using the ready-to-use
dye composition as defined above.

35 According to this process, at least one ready-
to-use dye composition as defined above is applied to

the fibres, for a period which is sufficient to develop the desired coloration, after which the fibres are rinsed, optionally washed with shampoo, rinsed again and dried.

5 The time required to develop the coloration on the keratin fibres is generally between 3 and 60 minutes and even more precisely between 5 and 40 minutes.

10 According to one specific embodiment of the invention, the process includes a first step which consists in separately storing, on the one hand, a composition (A) comprising, in a medium which is suitable for dyeing, at least one oxidation base and optionally at least one coupler as defined above, and,
15 on the other hand, a composition (B) containing, in a medium which is suitable for dyeing, at least one enzyme of 2-electron oxidoreductase type in the presence of at least one donor for the said enzyme and at least one aminosilicone, and then in mixing them
20 together at the time of use, before applying this mixture to the keratin fibres.

 According to another specific embodiment of the invention, the aminosilicone is incorporated into composition (A).

25 Another subject of the invention is a multi-compartment dyeing device or "kit" or any other multi-compartment packaging system, a first compartment of which contains composition (A) as defined above and a second compartment of which contains composition (B) as
30 defined above. These devices can be equipped with means for applying the desired mixture to the hair, such as the devices described in patent FR-2,586,913 in the name of the Applicant.

35 A subject of the present invention is also a novel process for treating keratin substances, in particular the hair, in order to obtain a permanent

0083207 044304

reshaping of this hair, in particular in the form of permanent-waved hair, this process comprising the following steps: (i) a reducing composition is applied to the keratin substance to be treated, the keratin substance being placed under mechanical tension before, during or after the said application, (ii) the keratin substance is optionally rinsed, (iii) an oxidizing composition as defined above is applied to the optionally rinsed keratin substance, (iv) the keratin substance is optionally rinsed again.

The first step (i) of this process consists in applying a reducing composition to the hair. This application is carried out lock by lock or all at once.

The reducing composition comprises, for example, at least one reducing agent, which can be chosen in particular from thioglycolic acid, cysteine, cysteamine, glyceryl thioglycolate, thiolactic acid or thiolactic or thioglycolic acid salts.

The usual step for placing the hair under tension in a shape corresponding to the desired final shape for this hair (for example curls) can be carried out by any suitable means, in particular mechanical means, which are suitable and known per se for maintaining the hair under tension, such as, for example, rollers, curlers and the like.

The hair can also be shaped without the aid of external means, simply with the fingers.

Before carrying out the following optional rinsing step (ii), the hair onto which the reducing composition has been applied should, conventionally, be left to stand for a few minutes, generally between 5 minutes and one hour, preferably between 10 and 30 minutes, so as to give the reducing agent enough time to act correctly on the hair. This waiting phase preferably takes place at a temperature ranging from 35°C to 45°C, while preferably also protecting the hair

with a hood.

In the optional second step of the process (step (ii)), the hair impregnated with the reducing composition is then rinsed thoroughly with an aqueous composition.

Next, in a third step (step (iii)), the oxidizing composition of the invention is applied to the hair thus rinsed, with the aim of fixing the new shape given to the hair.

As in the case of the application of the reducing composition, the hair onto which the oxidizing composition has been applied is then, conventionally, left for a standing or waiting phase lasting a few minutes, generally between 3 and 30 minutes, preferably between 5 and 15 minutes.

If the hair was maintained under tension by external means, these means (rollers, curlers or the like) can be removed from the hair before or after the fixing step.

Lastly, in the final step of the process according to the invention (step (iv)), which is also optional, the hair impregnated with the oxidizing composition is rinsed thoroughly, generally with water.

Hair which is soft and easy to disentangle is finally obtained. The hair is wavy.

The oxidizing composition according to the invention can also be used in a process for bleaching keratin fibres, and in particular the hair.

The bleaching process according to the invention comprises a step of applying an oxidizing composition according to the invention to the keratin fibres in the presence or absence of an auxiliary oxidizing agent. Conventionally, a second step of the bleaching process according to the invention is a step of rinsing the keratin fibres.

The medium which is suitable for the keratin


fibres (or the support) for the ready-to-use dye compositions and for the oxidizing compositions used for the permanent reshaping or bleaching of keratin fibres in accordance with the invention generally consists of water or of a mixture of water and at least one organic solvent in order to dissolve the compounds which would not be sufficiently soluble in water. By way of organic solvent, mention may be made, for example, of C₁-C₄ alkanols such as ethanol and isopropanol; glycerol; glycols and glycol ethers such as 2-butoxyethanol, propylene glycol, propylene glycol monomethyl ether, diethylene glycol monoethyl ether and monomethyl ether, and aromatic alcohols such as benzyl alcohol or phenoxyethanol, similar products and mixtures thereof.

The solvents can be present in proportions preferably of between 1 and 40% by weight approximately relative to the total weight of the dye composition, and even more preferably between 5 and 30% by weight approximately.

The pH of the ready-to-use dye compositions and of the oxidizing compositions used for the permanent reshaping or bleaching of the keratin fibres in accordance with the invention is chosen such that the enzymatic activity of the 2-electron oxidoreductase is not adversely affected. It is generally between 5 and 11 approximately, and preferably between 6.5 and 10 approximately. It can be adjusted to the desired value using acidifying or basifying agents usually used for dyeing keratin fibres.

Among the acidifying agents, mention may be made, by way of example, of inorganic or organic acids such as hydrochloric acid, orthophosphoric acid, sulphuric acid, carboxylic acids such as acetic acid, tartaric acid, citric acid or lactic acid, and sulphonic acids.

Among the basifying agents, mention may be made, by way of example, of aqueous ammonia, alkaline carbonates, alkanolamines such as mono-, di- and triethanolamines, 2-methyl-2-aminopropanol and derivatives thereof, sodium hydroxide, potassium hydroxide and the compounds of formula (IX) below:



10 in_ which W is a propylene residue optionally substituted with a hydroxyl group or a C₁-C₄ alkyl radical; R₂₂, R₂₃, R₂₄ and R₂₅, which may be identical or different, represent a hydrogen atom or a C₁-C₄ alkyl or C₁-C₄ hydroxyalkyl radical.

15 The ready-to-use dye compositions and the oxidizing compositions for the permanent reshaping or bleaching of keratin fibres in accordance with the invention can also contain various adjuvants used conventionally in compositions for dyeing, permanently
20 reshaping or bleaching the hair, such as anionic, cationic, nonionic, amphoteric or zwitterionic surfactants or mixtures thereof, anionic, cationic, nonionic, amphoteric or zwitterionic polymers or mixtures thereof, inorganic or organic thickeners,
25 antioxidants, enzymes other than the 2-electron oxidoreductases used in accordance with the invention, such as, for example, peroxidases, penetration agents, sequestering agents, fragrances, buffers, dispersing agents, conditioners, film-forming agents, preserving
30 agents and opacifiers.

Needless to say, a person skilled in the art will take care to select this or these optional complementary compound(s) such that the advantageous

00832877-041204

properties intrinsically associated with the compositions in accordance with the invention are not, or are not substantially, adversely affected by the addition or additions envisaged.

5 The ready-to-use dye compositions and the oxidizing compositions used for the permanent reshaping or bleaching of keratin fibres in accordance with the invention can be in various forms, such as in the form of liquids, creams or gels, which are optionally
10 pressurized, or in any other form which is suitable for dyeing, permanently reshaping or bleaching keratin fibres, and in particular human hair.

 In the case of a ready-to-use dye composition, the oxidation dyes(s) and the 2-electron oxido-
15 reductase(s) are present in the said composition, which must be free of oxygen gas, so as to avoid any premature oxidation of the oxidation dye(s).

Concrete examples illustrating the invention will now be given.

20 In the text hereinabove and hereinbelow, except where otherwise mentioned, the percentages are expressed on a weight basis.

 The examples which follow illustrate the invention without being limiting in nature.

25

0983287 041304
402719 4282860

Example 1 of dye composition:

The ready-to-use dye composition below was prepared (contents in grams):

5

- Uricase from *Arthrobacter globiformis* at a concentration of 20 International Units (I.U.)/mg, sold by the company Sigma 1.5 g
- Uric acid 1.5 g
- para-Phenylenediamine 0.324 g
- Resorcinol 0.33 g
- Polydimethylsiloxane containing amino-ethylaminopropyl groups, at a concentration of 35% in water, sold under the name Dow Corning 939 Emulsion 1.2 g
- Demineralized water qs 100 g

10 The ready-to-use dye composition described above was applied to locks of natural grey hair containing 90% white hairs for 30 minutes. The hair was then rinsed, washed with a standard shampoo and then dried.

Locks of hair dyed a matt dark-blonde colour were obtained.

Example 2: Oxidizing composition for permanent-
waving or bleaching

- Uricase from *Arthrobacter globiformis* at a
concentration of 20 International Units
(I.U.)/mg, sold by the company Sigma 1.8 g
- Uric acid 1.65 g
- Ethanol 20.0 g
- (C₈-C₁₀)alkyl polyglucoside as an aqueous
solution containing 60% active material
(A.M.), sold under the name Oramix CG110 by 8.0 g
the company SEPPIC
- Mixture of polydimethylsiloxane containing
aminoethylaminoisobutyl/polydimethyl-
siloxane groups, sold under the trade name
Dow Corning Q2 8220 Fluid 1.0 g
- 2-Methy-2-amino-1-propanol qs pH 9.5
- Demineralized water qs 100 g

00033297-04301